

## Multicomponent synthesis of 1, 2, 4-triazole-dihydropyrimidinone derivatives using Biginelli cycloaddition reaction

Kabeer Shaikh<sup>1</sup>, Yogesh Salve<sup>2\*</sup>

<sup>1</sup> Department of Chemistry, Sir Sayyed College of Arts, Commerce and Science College, Roshangate, Aurangabad, India.

<sup>2</sup> Department of Chemistry, Arts, Commerce and Science College Sonai, Tal- Newasa, Dist-Ahmednagar, India

### Abstract

An operationally simple protocol was employed for synthesis of series of new 1, 2, 4-triazole-dihydropyrimidinone derivatives. Firstly 1-(4-(1H-1, 2, 4-triazol-1-yl) phenyl) ethanone was synthesized using 4-fluoroacetophenone, 1, 2, 4-triazole, an.  $K_2CO_3$  in DMF solvent for 12 hrs at  $110^\circ C$ . then 1-(4-(1H-1,2,4-triazol-1-yl)phenyl) ethanone was refluxed with dimethyl foramide dimethylacetal (DMF-DMA) for 6 hrs gives product 1-(4-(1H-1,2,4-triazol-1-yl)phenyl)-3-(dimethylamino)prop-2-en-1-one (enaminone). Lastly different substituted aromatic aldehyde, urea and 1-(4-(1H-1, 2, 4-triazol-1-yl) phenyl)-3-(dimethylamino) prop-2-en-1-one (enaminone) was refluxed in glacial acetic acid for 12 hrs yields series of new 1,2,4-triazole-dihydropyrimidinone derivatives. The synthesized compound were characterized by mass,  $^1H$  NMR and  $^{13}C$  NMR spectroscopy followed by antibacterial and antifungal activity.

**Keywords:** 1, 2, 4-triazole, DMF-DMA, aromatic aldehyde, urea

### Introduction

Heterocyclic compounds containing bridgehead nitrogen are imperative for survival as they are part of crucial building blocks like amino acids, nucleotides, many coenzymes, metabolic regulators etc. Triazoles are one of the most essential nitrogen containing five membered heterocycles containing two carbon and three nitrogen atoms. There are two isomeric forms namely 1, 2, 3-triazole and 1, 2, 4-triazole <sup>[1,2]</sup> (fig.1).





Fig 1

The properties of triazoles have been utilized in many directions for years. The important structural accent of triazole ring with desirable electron feature is beneficial for triazole derivatives for easily binding of variety of enzymes and receptors in biological scenario to show activity against various diseases. They also having broad range of applications in pharmaceutical, agrochemical, supramolecular chemistry, organic synthesis<sup>3-6</sup>. Triazole and their fused heterocyclic compound derivatives serve interesting biological activities such as antibacterial<sup>7</sup>, antifungal<sup>8</sup>, antihypertensive<sup>9</sup>, anti-inflammatory<sup>10</sup> anticancer<sup>11</sup>, antitumor<sup>12</sup>, antimalarial<sup>13</sup>, antitubercular<sup>14</sup>, analgesic <sup>[15]</sup>. Triazole derivatives had shown potent antiviral activities against H1N1 influenza virus <sup>[16]</sup>. Among

derivatives of triazole, have been used in medical field as an antifungal agent such as Fluconazole, Itraconazole, Voriconazole, Triazolam, Furaciline, Alprazolam, Estazolam etc <sup>[17]</sup>. This disease is especially dangerous for people with minimized immunity, malnutrition, the elderly and alcohol abusers.



**Fig 2**

On the other side, dihydropyrimidinone also serve as interesting and multifaceted pharmacological activities such as antiviral <sup>[18]</sup>, antitumor <sup>[19]</sup>, antidiabetic <sup>[20]</sup>, antifungal and antibacterial <sup>[21]</sup>. As a consequence, dihydropyrimidinone derivatives synthesis containing contradictory substitution patterns has attracted marked attention as its discovery 120 years ago in 1893 by the Italian chemist Pietro Biginelli <sup>[22]</sup>. Among them, the Biginelli multicomponent reaction, involving a multicomponent condensation of aldehyde,  $\beta$ -ketoester, and urea in presence of different acidic or basic catalyst. By taking advantage from literature survey for above both (triazole and dihydropyrimidinone) bioactive heterocyclic moiety, most of researcher design and synthesized scaffolds exhibit significant biological activity. Guillermo *et al.* developed 1,2,3-triazole derivatives of dihydropyrimidinone as acidic corrosion inhibitor for steel <sup>[22]</sup>, Hassen *et al.* design and synthesized dihydropyrimidinone/1,2,3-triazole hybrid molecule as an antiviral agent <sup>[23]</sup>, Elisabete *et al.* design and synthesized a Novel 1,2,3-triazole-dihydropyrimidinone hybrids and evaluated anticancer activity <sup>[24]</sup>. Synthesis of Dihydropyrimidinone derivaives with imidazole moiety were synthesized by Biginelli reaction <sup>[25]</sup>. By considering all these above facts and literature survey suggested that compounds including these two important scaffolds (dihydropyrimidinone and triazole) should possess pharmaceutical activity so the target of present work is to synthesize new compounds containing dihydropyrimidinone derivatives with 1,2,4-triazole moiety.

### Experimental


All the chemicals were purchased from local suppliers and were used directly as received without further purification. The progress of reaction was monitored by Thin layer chromatography was accomplished on precoated plates of TLC silica gel 60 F<sub>254</sub>. Visualization was made with UV light (254 or 365nm). The mobile phase used for developing chromatogram is Methanol: DCM (1:10). The <sup>1</sup>H NMR spectra were recorded on BRUKER- 400 MHz (Model: AV 400) spectrometer in DMSO-d<sub>6</sub> solvent using TMS as an internal reference and Chemical shifts were supposed to be due to the presence of the solvent. Mass spectra were recorded on MS8040 Mass Spectrometer.

**Synthesis of (4-(1H-1, 2, 4-triazol-1-yl) phenyl) ethanone:** A mixture of 4-fluoro acetophenone (0.047 mol) and 1,2,4-triazole (0.047 mol) was taken in presence of potassium carbonate (0.056 mol) in DMF solvent. The reaction mixture refluxed for 12 hours, then reaction mixture was left to cool slowly at room temperature followed by addition of water to reaction mixture and extracted with ethyl acetate (3x 50 ml). The organic layer

was dry over an.  $\text{Na}_2\text{SO}_4$  and was concentrated under vacuum to get product. The obtained product was recrystallized from absolute ethanol.

**Synthesis of (E)-1-(4-(1H-1,2,4-triazol-1-yl)phenyl)-3-(dimethylamino)prop-2-en-1-one:** A mixture of (E)-1-(4-(1H-1,2,4-triazol-1-yl)phenyl) ethanone (0.02 mol) and dimethyl formamide-dimethyl acetal (DMF-DMA) (0.02 mol) was refluxed for 10 hours under solvent free condition on heating mantle. After the completion of reaction, the reaction mixture was slowly cooled to room temperature. The precipitate was obtained. Diethyl ether was added to the precipitate, and filtration was performed under vacuum. The obtained product (E)-1-(4-(1H-1, 2, 4-triazol-1-yl) phenyl)-3-(dimethylamino) prop-2-en-1-one was recrystallized from absolute ethanol.

**General Synthesis of the Triazole derivatives of Dihydropyrimidinone:** A mixture of (E)-1-(4-(1H-1, 2, 4-triazol-1-yl)phenyl)-3-(dimethylamino)prop-2-en-1-one (0.002 mol), different substituted benzaldehydes (0.002 mol), urea (0.002 mol), and glacial acetic acid (10 mL) was refluxed for 3hours. The precipitates were obtained by pouring the reaction mixture into the ice cold water. The products were isolated by Buchner filtration under vacuum. The products were washed three times with cold water followed by hexane. The physicochemical properties of different triazole derivatives are shown in Table. 1. In the  $^1\text{H-NMR}$  spectra, the signals of the individual protons of the compounds were verified on the basis of multiplicity, chemical shifts, and coupling constant. Analytical and Spectral data for the compounds were in good agreement with the expected structures of the compounds.



**Fig 3**

#### Name of structure

- 5-[4-(1H-triazole-1-yl)benzoyl]-4-phenyl-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta=9.38$  (1H, s, triazole H), 8.27 (1H, s, triazole H), 7.11 (1H, s, =CH), 5.44 (1H, d,  $J=2.8\text{Hz}$ ), 7.20-7.96 (m, 9H, Ar-H), 9.42 (s, 1H, NH), 7.88 (s, 1H, NH). MS:  $m/z=346$
- 5-[4-(1H-triazole-1-yl)benzoyl]-4-(2-methoxyphenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta=9.37$  (1H, s, triazole H), 8.28 (1H, s, triazole H), 3.83 (3H, s,  $\text{OCH}_3$ ), 7.13 (1H, s, =CH), 5.71 (1H, d,  $J=2.4\text{Hz}$ ), 6.89-7.97 (m, 8H, Ar-H), 9.34 (s, 1H, NH), 7.43 (s, 1H, NH). MS:  $m/z=376$
- 5-[4-(1H-triazole-1-yl)benzoyl]-4-(4-chlorophenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta=9.38$  (1H, s, triazole H), 8.27 (1H, s, triazole H), 7.12 (1H, s, =CH), 5.44 (1H, d,  $J=2.8\text{Hz}$ ), 7.34-7.96 (m, 8H, Ar-H), 9.49 (s, 1H, NH), 7.91 (s, 1H, NH). MS:  $m/z=380$
- 5-[4-(1H-triazole-1-yl)benzoyl]-4-(2-nitrophenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta=9.37$  (1H, s, triazole H), 8.26 (1H, s, triazole H), 7.17 (1H, s, =CH), 6.12 (1H, d,  $J=2.0\text{Hz}$ ), 7.50-8.10 (m, 8H, Ar-H), 9.60 (s, 1H, NH), 7.90 (s, 1H, NH). MS:  $m/z=391$
- 5-[4-(1H-triazole-1-yl)benzoyl]-4-(3-methoxyphenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta=9.38$  (1H, s, triazole H), 8.27 (1H, s, triazole H), 3.74 (3H, s,  $\text{OCH}_3$ ), 7.11 (1H, s, =CH), 5.42 (1H, d,  $J=2.8\text{Hz}$ ), 6.83-8.13 (m, 8H, Ar-H), 9.42 (s, 1H, NH), 7.87 (s, 1H, NH). MS:  $m/z=376$

- f. 5-[4-(1H-triazole-1-yl)benzoyl]-4-(4-trifluoromethylphenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta$ =9.37 (1H, s, triazole H), 8.26 (1H, s, triazole H), 7.14 (1H, s, =CH), 5.52 (1H, d,  $J$ =2.8Hz), 7.56-8.12 (m, 8H, Ar-H), 9.52 (s, 1H, NH), 7.95 (s, 1H, NH). MS:  $m/z$ =414
- g. 5-[4-(1H-triazole-1-yl)benzoyl]-4-(4-trifluoromethoxyphenyl)-3,4-dihydropyrimidin-2(1H)-one:  $^1\text{H NMR}$ : 400 MHz, DMSO- $d_6$ ):  $\delta$ =9.38 (1H, s, triazole H), 8.27 (1H, s, triazole H), 7.14 (1H, s, =CH), 5.48 (1H, d,  $J$ =2.8Hz), 7.23-7.96 (m, 8H, Ar-H), 9.50 (s, 1H, NH), 7.93 (s, 1H, NH). MS:  $m/z$ =430 Table.1: Physicochemical properties of compounds:

Table 1

Compound	R	Yield (%)
1	$\text{C}_6\text{H}_5$	70
2	2-OCH <sub>3</sub> - $\text{C}_6\text{H}_4$	68
3	4-Cl- $\text{C}_6\text{H}_4$	75
4	4-NO <sub>2</sub> - $\text{C}_6\text{H}_4$	72
5	3-OCH <sub>3</sub> - $\text{C}_6\text{H}_4$	68
6	2-NO <sub>2</sub> - $\text{C}_6\text{H}_4$	75
7	4-CF <sub>3</sub> - $\text{C}_6\text{H}_4$	68
8	4-OCF <sub>3</sub> - $\text{C}_6\text{H}_4$	68
9	4-F- $\text{C}_6\text{H}_4$	65
10	4-OCH <sub>3</sub> - $\text{C}_6\text{H}_4$	70




Fig 4

## Results and Discussion

As shown in scheme 1, reaction of 4-fluoro acetophenone with 1, 2, 4-triazole in presence of potassium carbonate and DMF solvent for refluxing 12 hrs yields (4-(1H-1, 2, 4-triazol-1-yl) phenyl) ethanone (III). (E)-1-(4-(1H-1, 2, 4-triazol-1-yl) phenyl)-3-(dimethylamino) prop-2-en-1-one (IV) was synthesized by refluxing (4-(1H-1, 2, 4-triazol-1-yl) phenyl) ethanone (III) with Dimethyl formamide-Dimethylacetal (DMF-DMA) under solvent free condition for 10 hrs. To prepare final triazole- dihydro pyrimidinone derivatives, a mixture of substituted aromatic aldehyde (V), enaminone (IV) and urea (VI) in glacial acetic acid was refluxed for 3 hrs. ) The ethylenic protons indicate that the enaminone existed in the E- configuration <sup>[26]</sup>. Compounds shows broad spectrum and having  $\delta$  value in between 7.43-7.95 ppm and 9.34-9.60 ppm indicates presence of two NH protons. The racemic proton of dihydropyrimidinone moiety is observed at 7.11-7.44 ppm  $\delta$  value as doublet ( $J=2.8\text{Hz}$ ). The  $\delta$  value in between 7.11-7.17 ppm indicates presence of allylic proton which appears as singlet. <sup>13</sup>C NMR and mass spectra of some compounds confirm the presence of compound.

The possible reaction mechanism for dihydropyrimidinone derivative containing 1, 2, 4-triazole (1–10) involves the acid-catalyzed formation of iminium ion intermediate from the substituted aryl aldehydes and urea. Reaction of iminium ion by enaminone of 1, 2, 4-triazole produces ureidenone, which cyclizes to form hexahydropyrimidine. Removal of N (CH<sub>3</sub>)<sub>2</sub> group from hexahydropyrimidine in presence of glacial acetic acid produces final dihydropyrimidinone derivative (1–10) containing 1, 2, 4-triazole moiety (Scheme 2)

## Conclusions

In conclusion, a series of dihydropyrimidinone derivatives with 1,2,4-triazole derivatives were synthesized with simple and efficient method having high purity of compound. The starting material was synthesized by 4-fluoro acetophenone with 1,2,4-triazole in presence of potassium carbonate in DMF solvent for 12 hrs which gives product acetophenone triazole compound. The enaminone were synthesized by treating acetophenone triazole with Dimethylformamide Dimethylacetal (DMF-DMA) under solvent free condition. The new dihydropyrimidinone derivatives were obtained from reaction of enaminone, substituted aryl aldehyde and urea in presence of glacial acetic acid.

## Acknowledgements

Authors are thankful to SAIF, Indian Institute of Science Bangalore for providing NMR facilities. Authors are grateful to Department of chemistry, Rashtrasant Tukadoji Maharaj Nagpur University, Nagpur for providing Mass facilities. Authors are sincerely appreciate to Principal, Arts, Commerce and Science College Sonai, Tal-Newasa, Dist-Ahmednagar, Maharashtra (India) for providing necessary facilities to carry out present work.

## References

1. Nadeem S, Waquar A, M Shamsheer A, Ruhi A, Sanjay J, Bishmillah A *et al.* Triazoles as potential bioactive agents. *Inter. J. Pharma. Sci. Review Research*,2011;8(1):61-169.[https://www.researchgate.net/publication/281875010\\_Triazoles\\_As\\_potential\\_bioactive\\_agents](https://www.researchgate.net/publication/281875010_Triazoles_As_potential_bioactive_agents)
2. Jawad KS, Yusra HA. Chemistry of 1, 2, 4-triazole: a review article. *Inter. J. Sci. and Research*,2016;5(3):1411-1423.[https://www.researchgate.net/publication/312021996\\_Chemistry\\_of\\_1\\_2\\_4-Triazole\\_A\\_Review\\_Article](https://www.researchgate.net/publication/312021996_Chemistry_of_1_2_4-Triazole_A_Review_Article)
3. Gang Y, Serajul H, Li S, Gnanasambandam K, Joy W, Thomas Me *et al.* Synthesis of alkyne derivatives of a novel triazolopyrazine as A<sub>2</sub>A adenosine receptor antagonists. *Bioorg. Med. Chem. Lett*,2005;15:511-515. <https://doi.org/10.1016/j.bmcl.2004.11.062>
4. Wei H, Guang-fu Y. Microwave-assisted, one-pot syntheses and fungicidal activity of polyfluorinated 2-benzylthiobenzothiazoles. *Bioorg. Med. Chem. Lett.* 14, 2006, 8280-8285.DOI: 10.1016/j.bmc.2006.09.016
5. Benjamin S, Ulrich SS. Beyond click chemistry – supramolecular interactions of 1, 2,3-triazoles. *Chem. Soc. Rev.*, 43, 2014. 2522-2571.DOI: <https://doi.org/10.1039/c3cs60386e>
6. Ahmed AE, Soliman NK. Synthesis, biological activity and molecular modeling study of novel 1,2,4-triazolo[4,3-*b*][1,2,4,5] tetrazines and 1,2,4-triazolo[4,3-*b*] [1,2,4]triazines *Scientific Reports*, 10, 2020, 6137-6155.DOI:10.1038/s41598-020-62977-x
7. Bo Z, Comprehensive review on the anti-bacterial activity of 1,2,3-triazole hybrids. *Eur. J. Med. Chem.*, 168, 2019, 357-372.DOI: 10.1016/j.ejmech.2019.02.055.
8. Davir GC, Maria GM, Marco AM, Alejandra RV, Macario MR, Bertha JR *et al.*, Aydee FB, Azide-enolate 1,3-dipolar cycloaddition in the synthesis of novel triazole-based miconazole analogues as promising antifungal agents. *Eur. J. Med. Chem.*,112, 2016, 60-65.<https://doi.org/10.1016/j.ejmech.2016.02.013>
9. Jie L, Qin L, Xue Y, Shengtao X, Hengyuan Z, Renren B *et al.* Design, synthesis, and biological evaluation of 1,2,4-triazole bearing 5-substituted biphenyl-2-sulfonamide derivatives as potential antihypertensive candidates. *Bioorg. Med. Chem.* 21, 2013, 7742-7751. DOI: 10.1016/j.bmc.2013.10.017
10. Renata P, Malgorzata W, Andrzej E, Anna HB, Andrzej G, Bozena MB *et al.* Synthesis and anti-inflammatory activity of new 1,2,4-triazole activity, *Bioorg. Med. Chem. Lett.* 25, 2015, 2664-2667.DOI: 10.1016/j.bmcl.2015.04.079
11. Yazala JP, Reddymasu S, Deekala V, Rudraraju RR. Design, Synthesis, and Biological Evaluation of 1, 2, 4-Thiadiazole-1, 2, 4-Triazole Derivatives Bearing Amide Functionality as Anticancer Agents. *Arabian J. Sci. Engi.*, 46, 2021, 225-232. <https://doi.org/10.1007/s13369-020-04626-z>

12. Harbinder S, Mandeep K, Kunal N, Manish KG, Ajit KS, Sahil S *et al.* Triazole tethered C5-curcuminoid-coumarin based molecular hybrids as novel antitubulin agents: Design, synthesis, biological investigation and docking studies. *Euro. J. Med. Chem.*, 116, 2016, 102-115. DOI: 10.1016/j.ejmech.2016.03.050
13. Kewal K, Bruno P, Marilyn M, Remy A, Nicolas B, Vipin K. 1H-1,2,3-triazole tethered isatin-ferrocene conjugates: Synthesis and in vitro antimalarial evaluation. *Euro. J. Med. Chem.*, 2014;87:801-804. <http://dx.doi.org/10.1016/j.ejmech.2014.10.024>
14. Raggappa SK, Siddapa AP, Srinivasa B, Bhari MN. Triazole: A Promising Antitubercular Agent. *Chem Biol Drug Des*, 2015;86(4):410-423. DOI: 10.1111/cbdd.12527
15. Umut SG, Nesrin GK, Ozgur G, Yavuz K, Ekrem K, Samil I *et al*, Meral O. 1-Acylthiosemicarbazides, 1,2,4-triazole-5(4H)-thiones, 1,3,4-thiadiazoles and hydrazones containing 5-methyl-2-benzoxazolinones: Synthesis, analgesic-anti-inflammatory and antimicrobial activities. *Bioorg. Med. Chem*, 2007;15:5738-5751. DOI: 10.1016/j.bmc.2007.06.006
16. Nashwa HZ, Mohammed IM, Abdullah YA. Design, synthesis and molecular docking of novel triazole derivatives as potential CoV helicase inhibitors. *Acta Pharm*, 2020;70:145-159. <https://doi.org/10.2478/acph-2020-0024>
17. Armen SG, Tariel VG, Melanya AS, Samvelyan VR, Frangyan M Sarfraz. *Chemistry Select*, 2019;4:12386-12390. DOI: 10.1002/slct.201902761
18. Nicholas P, Peter T, Margrith EM, Li Y, Beth AS, Zhaohui C *et al.* An integrated chemical biology approach reveals the mechanism of action of HIV replicaion inhibitors. *Bioorg. Med. Chem*, 2017;25:6248-6265. <http://dx.doi.org/10.1016/j.bmc.2017.03.061>
19. Mariana M, Goncalo C, Adriana OS, Amilcar F, Samuel S, Gilberto A. Potential aniumoral 3,4-dihydropyrimidin-2-(1H)-ones: synthesis, in vitro biological evaluation and QSAR studies. *RSC Advnaces*, 2016;88, 84943-84958. <https://doi.org/10.1039/C6RA14596E>
20. Kashinath LD, Surya NM, Sanjeev CG, Sanosh GT. Graphite catalyzed solvent free synthesis of dihydropyrimidin-2(1H)-ones/thiones and their anti-diabetic activity. *Bioorg. Medi. Chem. Lett*, 2014;24:2897-2901. <http://dx.doi.org/10.1016/j.bmcl.2014.04.099>
21. Tarunkumar NA, Jignesh PR. 1,3-dihydro-2H-indol-2-ones derivatives: Design, Synthesis, in vitro antibacterial, antifungal and antitubercular study. *Eur. J. Med. Chem*, 2011;46:5573-5579. DOI: 10.1016/j.ejmech.2011.09.023
22. Rodrigo GO, Viridiana RR, Guillermo EN, Araceli EV, Francisco JR Rosa S. Multicomponent synthesis and evaluation of new 1,2,3-triazole derivatives of dihydropyrimidinones as acidic corrosion inhibitors for steel. *Molecules*, 2016;21:250-263. DOI: 10.3390/molecules21020250
23. Hanane K, Youssef K, Christophe C, Moha T, Juan CR, Robert S, Graciela A, Patrice V, Hassan BL. Dihydropyrimidinone/1,2,3-triazole hybrid molecules: synthesis and anti-varicella-zoster virus (VZV) evaluation. *Eur. J. Med. Chem.* 155, 2018, 772-781. DOI: 10.1016/j.ejmech.2018.06.028
24. Elisabete PC, Ana MS, Adrian P, Jose MP, Anthony JB. Synthesis of novel 1, 2, 3-triazole-dihydropyrimidinone hybrids using multicomponent 1, 3-dipolar cycloaddition (click)-biginelli reactions: Anticancer activity. *Synlett*, 31, 2020, 615-622. DOI: 10.1055/s-0039-1690781; Art ID: st-2019-b0556-1
25. Mashooq AB, Mohamed AA, Ahmed MN, Atef K, Abdullah AD. *J. Chem.* 2019. <https://doi.org/10.1155/2019/3131879>
26. Mashooq AB, Atallah FA, Zhi W, Mohamed AA Hahem AA. *Med Chem Res*, 2017;26:1557-1567. DOI: 10.1007/s00044-017-1870-5