

BIOACCUMULATION OF HEAVY METALS IN TISSUES OF *LABEO ROHITA* FROM MULA DAM, AT RAHURI, AHMEDNAGAR, INDIA

M. J. SHAIKH

Department of Zoology, Arts, Commerce & Science College,
Sonai, Maharashtra, India,
e-mail- mjshaikh1850@gmail.com

ABSTRACT

The present study was conducted to determine the bioaccumulation of heavy metals (Cadmium and Chromium) in tissues (Gills and Liver) of the fish *Labeo rohita* from Mula Dam, near Rahuri in Maharashtra. The concentration of cadmium in sample water collected from site I was maximum 0.0075 mg/l, which is more than permissible limits for drinking water. The concentration of chromium in sample water is however less and the mean was 0.059 mg/l. The accumulation of cadmium in liver was found more (1.83 mg/kg) compared to gills (1.62 mg/kg). The maximum bioaccumulation factor (BAF) of cadmium was 280.00 in the liver and maximum BAF of cadmium was 241.79 in the gills. The concentration of chromium was found more (3.14 mg/kg) in liver compared to gills (2.81 mg/kg). The BAF of chromium was found maximum 53.10 in liver and 47.06 in gills. However, the accumulation of cadmium in both the tissues of *Labeo rohita* was less than permissible limits, but that of chromium was found to be more.

Figure: 00

References: 09

Tables: 03

KEY WORDS: *Labeo rohita*, Mula Dam, Cadmium, Chromium, Bioaccumulation factor.

Introduction

In aquatic ecosystem, heavy metals are considered as the most important pollutants, since they are present throughout the ecosystem and are detectable in critical amounts. Heavy metals are non-biodegradable and once discharged into water bodies, they can either be adsorbed on sediment particles or accumulated in aquatic organisms. Fish may absorb dissolved elements and heavy metals from surrounding water and food, which may accumulate in various tissues in significant amounts and shows toxicological effects at critical targets [3].

The bioaccumulation of heavy metals in different fish tissues has been studied by several investigators [4],[5]. Fish is often at the top of aquatic food chain and may concentrate large amounts of some metals from the water. Metal bioaccumulation is largely attributed to differences in uptake and depuration period of various metals in different fish species. Multiple factors including season, physical and chemical properties of water can play a significant role in

metal accumulation in different fish tissues. The gills are directly in contact with water. Therefore, the concentration of metals in gills reflects their concentration in water where the fish lives, whereas the concentration in liver represents storage of metals in water [2].

The heavy metals, being conservative in nature have the maximum probability of biomagnifications, when they are transferred to the human beings through the various members of different tropic levels in the food chain. Human beings are affected negatively as a result of their accumulation. In the Mula Dam, Maharashtra, India there is ample of fish *Labeo rohita* (Rohu) and commonly used as aquatic food by the people. This study aims to find out the bioaccumulation of heavy metals (Cadmium and Chromium) in different tissues (gills and liver) of the fish *Labeo rohita* collected from three different sites of Mula Dam.

Materials and Methods

Water samples and fish samples (*Labeo rohita*) were collected from three different experiments were analyzed and the results were expressed as mean of all three. Values of $P < 0.05$ were considered statistically significant.

Bioaccumulation factor (BAF) is the ratio between the accumulated concentration of a given pollutant in any organ and its dissolved concentration in water. It is calculated by dividing concentration of heavy metals in fish organs by heavy metals found in water. sites of Mula Dam, near Rahuri, Ahmednagar in December 2015. Site I was near the wall of Dam and sites II and III were 500 meters distance away respectively from site I. Water samples were collected at 50 cm below the surface, using 1 liter polythene bottle with screw caps. The bottle had been washed and soaked in 5% nitric acid and rinsed with deionised water before use. The water samples were acidified immediately after collection by adding 5 ml nitric acid to minimize adsorption of heavy metals onto the walls of the bottles [1]. Water samples were analyzed in UV Spectrophotometer for detection of heavy metals.

Samples of three fish (*Labeo rohita*) of nearly equal size and weight were dissected to remove muscles, gills and liver. The separated organs were put into petri dishes to dry at 120°C . The organs were placed into digestion flasks and ultrapure Con. Nitric acid and hydrogen peroxide (1:1 v/v) was added. The digestion flasks were then heated to 130°C until all the materials were dissolved. Digest was diluted with double distilled water appropriately. The heavy metals Cadmium (Cd) and Chromium (Cr) were assayed using UV Spectrophotometer and the results were given as mg/kg dry weight. Data obtained from the experiments were analyzed and the results were expressed as mean of all three. Values of $P < 0.05$ were considered statistically significant.

Bioaccumulation factor (BAF) is the ratio between the accumulated concentration of a given pollutant in any organ and its dissolved concentration in water. It is calculated by

dividing concentration of heavy metals in fish organ by heavy metals found in water.

Result and Discussion

Heavy metal concentrations in the sample water collected from three different sites of Mula Dam is given in Table - 1. There is fluctuation in the concentration of Cadmium (Cd) as well as Chromium (Cr). Maximum amount of Cd found is 0.0075 mg/l at site I, near the Dam. The regional mean of Cd was observed to be 0.0069 mg/l which is more than permissible limit of WHO, 2004 for drinking water. The concentration of Cr found is 0.061 mg/l at site III. The regional mean of Cr dissolved in sample water was 0.059 mg/l, which is more than permissible limit (0.05 mg/l) of WHO for drinking water.

The concentration of Cd in gills and liver of fish *Labeo rohita* is given in Table-2. The accumulation of Cd is more in liver tissues than gills. The liver shows maximum (1.83 mg/kg dw) amount of Cd, where as gills show 1.62 mg/kg as the maximum concentration. The regional mean of Cd found in gills is 1.57 mg/kg where as that of liver is 1.79 mg/kg. The values of Cd found both in gills and liver is lower than permissible limits of WHO.

The concentration of Cr in gills and liver of fish *Labeo rohita* is given in Table-3. The accumulation of Cr is more in liver tissues than gills. The liver shows maximum 3.14 mg/kg of Cr, where as gills show 2.81 as the maximum concentration. The regional mean of Cr found in gills is 2.74 where as that found in liver is 3.11. The values of Cr observed in gills and liver exceeds the permissible limits of WHO. Similar results are observed by [8] in the organs of fish *Cyprinus carpio* from Tamilnadu and by [7] working on River Yamuna in Delhi.

It was found that the concentration of Cd and Cr in both the tissues of fish were several times higher than their concentrations in water. The bioaccumulation factors (BAF) are given in Table-2 and Table-3. Cadmium shows highest BAF (280.00) in the liver and lowest (208.00) in gills. Comparatively, bioaccumulation of Chromium is lower than Cadmium. However, the lowest values of BAF

were recorded in gills. Bioaccumulation factor gives an indication about the accumulation efficiency for any particular pollutant in any fish organs. Similar results were observed by [6] in the fish *Tilapia nilotica*. The results suggested the loss of homeostatic capacity of fish under chronic metal exposure leading to bioaccumulation.

Conclusion

The results of this study supply valuable information on the metal contents in water sample and fish organs of the river Mula, at

TABLE 1: Heavy metal concentration (mg/l) in the water of Mula Dam, Rahuri

Sites	Cd	Cr
I	0.0075	0.061
II	0.0067	0.058
III	0.0065	0.059
Regional mean	0.0069	0.059
WHO limit for Drinking water	0.005	0.05

TABLE 2: Concentration of Cadmium (mg/kg dry weight) and the bioaccumulation factor (BAF) in different tissues of *Labeo rohita* from Mula Dam, Rahuri

Sites	Gills	Liver
I	1.56 (208.00)	1.72 (229.33)
II	1.62 (241.79)	1.83 (273.13)
III	1.53 (235.38)	1.82 (280.00)
Regional mean	1.51	1.79

Mula Dam. Fish liver exhibited highest tendency to accumulate both cadmium and chromium. The bioaccumulation of chromium in gills and liver is beyond the standard limits. The result indicates that the heavy metal contamination definitely affects the aquatic life of the fresh water fish. Hence, a scientific method detoxification is essential to improve the life of these economically important fishes in any stressed environmental conditions.

WHO limit for fish tissues	2.0	2.0
----------------------------	-----	-----

Values in brackets are bioaccumulation factors (BAF)

TABLE 3: Concentration of Chromium (mg/kg dry weight) and the bioaccumulation factor (BAF) in different tissues of *Labeo rohita* from Mula Dam, Rahuri

Sites	Gills	Liver
I	2.81 (46.07)	3.14 (51.47)
II	2.73 (47.06)	3.08 (53.10)
III	2.68 (45.42)	3.12 (52.88)
Regional mean	2.74	3.11
WHO limit for fish tissues	2.8	2.8

Values in brackets are bioaccumulation factors (BAF)

References

1. APHA-AWWA. (2004). Standard methods for examination of water and waste water. 21st Edition, Washington DC., USA.
2. RAUF, A., JAVED, M. AND UBaidULLAH, M. (2009). Heavy metal levels in three major carps (*Catla catla*, *Labeo rohita* and *Cirrhina mrigala*) from the river Ravi, Pakistan. *Pakistan Vet. J.*, 29(1): 24-26.
3. CALTA, M. AND O.CANPOLAT (2006). The comparison of three cyprinid species in terms of heavy metal accumulation in some tissues. *Water Environmental Research*, 78: 548 – 551.
4. FATMA, A.S., MOHAMED. (2008). Bioaccumulation of selected metals and histopathological alterations in tissues of *Oreochromis niloticus* and *Lates niloticus* from lake Nasser, Egypt. *Global Veterinaria*, 2 (4): 205 – 218.
5. JAVED, M. AND USMANIM, N. (2011). Accumulation of heavy metals in fishes : A human health concern. *International Journal of Environmental Sciences*, 2 (2): 659 – 668.

6. RASHED, M.N. (2001).Cadmium and lead levels in fish (*Tilapia nilotica*) tissues as biological indicator for lake water pollution. *Environmental Monitoring and Assessment*, **68**: 75-89.
7. SHRIVASTAVA,V.S.,AJAY, SHANDIL. AND INDRAJIT, SEN. (2011).Study of determination of heavy metals in fish species of the River Yamuna (Delhi) by Inductively Coupled Plasma Optical Emission Spectroscopy. *Adv. Appl. Sci. Res.*, **2** (2) : 161 – 166.
8. VINODHINI, R. AND NARAYANAN, M.(2008).Bioaccumulation of heavy metals in organs of fresh water fish *Cyprinus carpio* (Common carp). *Int. J. Environ. Sci. Tech.*, **5**(2): 179-182.
9. WHO.(2004)List of maximum levels recommended for contaminants by the Joint FAO/WHO Codex Alimentarius Commission.